

THIE UNITED STAYLES OF AMIERICA

TO ME TO WHOM THESE: PRESENTS SHALL COME:: California Cooperative Rice Research Joundation, Inc.

DUCCUS, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS EXOM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR FRERENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN UNITED STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. AMENDED, 7 U.S.C. 2321 ET SEQ.)

RICE

'M-206'

In Vestimon Musser, I have hereunto set my hand and caused the seal of the Mant Pariety Archertism Office to be affixed at the City of Washington, D.C. this tenth day of November, in the year two thousand and four.

De M J. __ Commissioner

Plant Variety Protection Office Agricultural Marketing Service Antenan

Secretary of Agriculture

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY - PLANT VARIETY PROTECTION OFFICE The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (FRA) of 1995. Application is required in order to determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential until certificate is issued (7 U.S.C. 2425). FLICATION FOR PLANT VARIETY PROTECTION CERTIFICATE (Instructions and information collection burden statement on reverse) AME OF OWNER 2. TEMPORARY DESIGNATION OR EXPERIMENTAL NAME 3. VARIETY NAME California Cooperative Rice Research Foundation, Inc. 98-Y-242 M - 206CORESS (Street and No., or R.F.D. No., City, State, and ZIP Gage, and Country) 5. TELEPHONE (Include area code) FOR OFFICIAL USE ONLY Rice Experiment Station PVPO NUMBER 530-868-5481 P.O. Box 306 955 Butte City Hwy 6. FAX (include area code) Biggs, CA 95917-0306 530-868-1730 FILING DATE THE OWNER NAMED IS NOT A "PERSON", GIVE FORM OF RGANIZATION (carporation, partnership, association, etc.) March 4, 2003 9. DATE OF INCORPORATION Non Profit Organization 1912 NAME AND ADDRESS OF OWNER REPRESENTATIVE(S) TO SERVE IN THIS APPLICATION. (First person listed will receive all papers) FILING AND EXAMINATION Dr. Kent S. McKenzie 3652.*0*0 Director DATE Feb. 20, 2003 Rice Experiment Station Biggs, CA 95917-0306 432.00 DATE 10-5-14 TELEPHONE (Include area code) 12. FAX (Include area code) 13. E-MAIL 14. CROP KIND (Common Name) 530-868-5481 530-868-1730 ksmckenzie@crrf.org Rice SENUS AND SPECIES NAME OF CROP 17. IS THE VARIETY A FIRST GENERATION HYBRID? 16. FAMILY NAME (Estanical) Oryza sativa L. Gramineae 🔁 мо ☐ YES. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE SOLD AS A CLASS OF CERTIFIED SEED? See Section 83(a) of the Plant Variety Protection Act) Exhibit A. Origin and Breeding History of the Variety 粒 YES (If "yes", answer items 20 and 21 below) a. X ☐ NO (If "no", go to item-22) o. 🔯 Exhibit 8. Statement of Distinctness Embit C. Objective Description of Variety 20. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF CLASSE X YES ☐ NO Exhibit O. Additional Description of the Variety (Optional) IFYES, WHICH CLASSES? Z FOUNDATION ★ CERTIFIED REGISTERED X Exhibit E. Salement of the Basis of the Owner's Ownership Voucher Sande (2,500 viable untreated seeds or, for tuber propagated varieties, verification that tissue culture will be deposited and maintained in an approved public. X 21. DOES THE OWNER SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF GENERATIONS? NΩ Filing and Examination Fee (\$2,705), made payable to "Treasurer of the United States" (Mail to the Plant Variety Protection Office) g_ 🔲 FYES: SPECIFY THE 2 FOUNDATION CERTIFIED 1 NUMBER 1.23 etc. 4,, (If additional explanation is necessary, please use the space indicated on the reverse.) HAS THE VARIETY (INCLUDING ANY HARVESTED MATERIAL) OR A HYBRID PRODUCED FROM THIS VARIETY BEEN SOLD, DISPOSED OF, TRANSFERRED, OR USED IN THE U.S. OR IS THE VARIETY OR ANY COMPONENT OF THE VARIETY PROTECTED BY INTELLECTUAL PROPERTY RIGHT (PLANT BREEDER'S RIGHT OR PATENT)? OTHER COUNTRIES? X YES ☐ NO XX YES IF YES, YOU MUST PROVIDE THE DATE OF FIRST SALE, DISPOSITION, TRANSFER, OR USE FOR EACH COUNTRY AND THE CIRCUMSTANCES. (Please use space indicated on reverse.) IFYES, PLEASE-GIVE COUNTRY, DATE OF FILING OR ISSUANCE AND ASSIGNED. REFERENCE NUMBER. (Flease use space indicated on reverse.) The owners declare that a visible sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a luber propagated watery a tissue culture will be deposited in a public repository and maintained for the duration of the certificate. The undersigned owners) is(are) the owner of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act. Owner(s) is(are) informed that false representation herein can jeopardize protection and result in penalties. SIGNATURE OF OWNER NATURE OF OWNER IE (Please print or type) NAME (Please print ontype) ACITY OR TITLE CATE CAPACITY OR TITLE 2-13-03 70 (04-01) designed by the Ffant Vaniery Protection Office with Worth Perfect 5.0a. Replaces STD-470 (02-99

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in the PVPC: (1) Completed application form signed by the owner, (2) completed exhibits A, B, C, E; (3) for a seed reproduced variety at least 2,500 viable untreated seeds, for a hybrid variety at least 2,500 untreated seeds of each line necessary to reproduce the variety, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture will be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,705 (\$320 filing fee and \$2,385 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice.) Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application and other requirements to Plant Variety Protection Office, AMS, USDA, Room 401, NAL Building, 10301 Baltimore Avenue, Beltsville, MD 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below. Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. If a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$320 for issuance of the certificate. Certificates will be issued to owner, not licensee or agent.

Plant Variety Protection Office Telephone: (301) 504-5518 FAX: (301) 504-5291

Homepage: http://www.ams.usda.gov/science/pvpo/pvp.htm

ITEM
18a. Give:

- (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
- (2) the details of subsequent stages of selection and multiplication;

(3) evidence of uniformity and stability; and

- (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified
- 18b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all other varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;

(2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences; and

- (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 18c. Exhibit C forms are available from the PVPO Office for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- 18d. Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 18e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 19. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant MAY NOT reverse this affirmative decision after the variety has been sold and so labeled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 22. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.
- 23. See Section 55 of the Act for instructions on claiming the benefit of an earlier filing date.
- 21. CONTINUED FROM FRONT (Please provide a statement as to the limitation and sequence of generations that may be certified poundation seed is maintained by the Rice Experiment Station(RES). Normal seed multiplication starts with head row seed used for breeders which in turn is used to plant foundation. Headrow seed is produced as necessary. RES policy allows foundation seed to be used to produce Foundation seed when necessary continued from the date of first sale, disposition, transfer, or use for each country and the circumstances, if the variety (including any harvested material) or a hybrid produced from this variety has been sold, disposed of, transferred, or used in the U.S. or other countries.) M-206 was jointly released by the California Co-op Rice Research Foundation, California Agricultural Experiment Station, and the USDA-ARS. Foundation will be distributed to California rice seed growers April 1, 2003 for the purpose of producing registered seed.
- 23. CONTINUED FROM FRONT (Please give the country, date of filing or issuance, and assigned reference number, if the variety or any component of the viety is protected by intellectual property right (Plant Breeder's Right or Patent).)

M-206 application Docket No. 1351-0008, Serial No. 10/657,892 issued 9 Sept.

RAO 914/04

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment or owner's representative during the life of the application/certificate. There is no charge for filling a change of address. The fee for filling a change of ownership or assignment or any modification of owner's name is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Sections 97.130, 97.131, 97.175(h) of the Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant must check the appropriate recognized authority. For example, for agricultural and vegetable crops, contact: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center—East, Beltsville, MD 20705. Telephone: (301) 504-8089. http://www.ams.usda.gov/lsg/seed.htm

According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and a person is not required to respond to a collection of information unless it displays a valid OMB control number. The valid OMB control number for this information collection is 0581-0055. The time required to complete this information collection is estimated to average 3.0 hours per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed; and completing and reviewing the collection of information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, sax, religion, age, disability, political beliefs, sexual crientation, or marital or family status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact. USDA's:TARGET Center at 202-720-2800 (voice and TDD).

To file a complaint of discrimination, write-USDA, Director, Office of Civil Rights, Room 328-W, Whitten Building, 14th and Independence Avenue, SW, Washington, DC 20250-9410 or call 202-720-5984 (voice and TDD). USDA is an equit opportunity provider and employer.

S&T-470 (07-01) designed by the Plant Variety Protection Office with WordPerfect 9.0. Replaces STD-470 (04-01) which is obsolete.

'M-206' Rice

18a. Exhibit A. Origin and Breeding History of the Variety

1) Genealogy

98-Y-242 is a very early, to early, semidwarf, California glabrous, Calrose quality medium-grain (CRMG) experimental line that has been evaluated in large plot yield tests since 1998. It is a very early selection from the 1991-92 winter greenhouse cross R18087 and has the pedigree S-301/M-204. Both varieties used in the pedigree were developed by the California Cooperative Rice Research Foundation, (CCRRF) at the Rice Experiment Station (RES), Biggs, CA. S-301 is an intermediate maturity short-grain semidwarf that is no longer in production. M-204 is an early maturing Calrose medium-grain with semidwarf height. 98-Y-242 is the product of modified pedigree breeding. Early spray painted panicle selections from the 1993 F₂ nursery selected for medium grain shape were bulked 6 panicles per row in the 1993-94 Hawaiian nursery. Panicle selections grouped from the R18087 Hawaii rows were planted in 1994 individual rows. The line of descent was from 94 row 63811 via individual panicle selection to 95 row 6309 to 96 row 5136 which was bulked for 97 small plot 351 that was the seed source for entry 98-Y-242 in the RES Preliminary Yield Test. It has been tested in the University of California Cooperative Extension (UCCE) Statewide Yield Tests (99-Y-84, 00-Y-63, 01-Y-12, 01-Y-72, 02-Y-11, 02-Y-73) 1999 to 2002.

2) Selection and multiplication

Selection was made for, improved lodging resistance, early maturity, cold tolerance and yield potential, and grain quality characteristics similar to Calrose medium grains. Panicle selections were first taken for headrow production in 1999 and continued through 2002. The 1999 panicles were initially purified in the 2000 headrow nursery. Year 2000 selected headrows were generation advanced in 2000-01 Hawaiian nursery. Year 2001 breeder headrows totaled 300 rows from 2000 selected headrows and their descendents from 2001-02 Hawaii nursery. The year 2001 breeder increase consisted of two components. One component was derived from bulked year 2000 headrows while the other came from the 2000-01 Hawaii nursery. These bulks were harvested separately and planted in different foundation field locations in 2002 for a total of 7.7 acres. Previous experience revealed that in certain years more outcrossing occurs in the Hawaii nursery resulting in more offtypes in either the breeder and/or foundation production fields. Two years of observations and plant counts demonstrated that 2000-01 outcrossing in Hawaii was near non-existent resulting in the two components being combined for 2002 Foundation seed. The 2001 headrow panicle selections were used to plant 400 headrows in 2002.

3) Uniformity and stability

Panicle selections were first taken for headrow production in 1999 and continued through 2002. During the headrow years any putative segregating or variant rows were removed from the field. Visual inspection (seedling vigor, heading date, plant height, grain shape and size, etc.) of headrows were used as criteria to confirm cultivar purity. This is standard procedure at RES to monitor and maintain breeder seed of the 13 currently grown

California rice varieties. M-206 was approved for certification by the California Crop Improvement Association in 2002. Classes of seed will be breeder, foundation, registered, and certified seed produced in California. Foundation seed can be used to produce foundation seed if necessary and headrow and breeder seed will be produced in foundation fields as necessary to maintain cultivar purity. M-206 has been observed in seed increase and production fields for four generations (2000 to 2002) and found to be uniform and stable.

4) Variants

The breeder, foundation, and headrows were rogued multiple times and off-types, putative out crosses, and other visual variants were removed. These included pubescent and glabrous longer medium-grains with and without red apiculus, pubscent shorter length medium-grain, slightly earlier and later maturing medium-grains, and fully awned medium grains possibly from outcrossing. In addition there were putative tetraploids (larger kernel size, delayed maturity and high floret sterility) and chlorophyll chimeras removed from the field. Total frequency of these off types was 0.0172%.

18b. Exhibit B. Statement of Distinctiveness

1) Variety Differences

M-206 is a semidwarf, very early to early maturing, medium-grain being released to serve the Calrose medium-grain market. "Calrose" was the founding California medium-grain rice variety, the ancestor of California medium-grains, and is now recognized as a market class term for California medium-grain rice. Comparisons for evaluation purposes were made to M-202, which is the predominant early maturing California medium grain. M-104 is a very early Calrose medium grain. M-206 maturity (days 50% flowering) averages 1/2 way between M-104 and M-202. M-206 has higher harvest moisture than M-104. M-206 is most similar to M-202 and is distinguished by its earlier heading date.

2) Statistical data

Agronomic characteristics: Agronomic characteristics (seedling vigor through yield) were collected in multiple locations (N=21) University of California Cooperative Extension (UCCE) Statewide Yield Tests in 1999 to 2002 are summarized in Table 1. M-206 is significantly later than M-104 and significantly earlier than M-202. M-206 has significantly higher harvest moisture than M-104. Tables 2 & 3 show individual year statistical evidence collected from replicated large plot yield tests at RES.

Table 1. Summary of Agronomic Characteristics for M-104, M-206, and M-202 from 1998 to 2002.

Character	M-104	M-206	M-202
Seedling vigor (score)	4.5	4.5	4.6
Days 50% heading	82	*85*	89
Plant height (cm)	93	95	. 97
Lodging (%)	50	31	38
Blanking – Greenhouse (%)	13	12	25
Blanking – Davis (%)	7	12	16
Blanking San Joaquin (%)	9	7	27
Overall blanking mean (%)	13	10	23
Stem rot (score)	5.6	5.4	5.5
Harvest moisture (%)	17.6	*19.8	19.7
Yield (lb/acre @ 14%)	9087	9350*	9060
Milling (%) total	68.9	69.0	68.7
Milling (%) whole grain	63.4	66.0	64.2
Brown weight 1000 seeds (g)	24.0	24.4	22.7
Brown length (mm)	6.26	6.23	5.96
Brown width (mm)	2.72	2.75	2.80
Ratio (L/W)	2.30:1	2.27:1	2.13:1
Apparent Amylose (%)	17.5	17.4	16.5
Alkali score (1.7% KOH)	6.5	6.8	6.9

^{*}Significantly different at the 0.05 probability level.

Table 2. Harvest moisture of M-104, M-206, and M-202 from 1999-2002 at the Rice Experiment Station, Biggs, CA.

Year	Trial	M-104		M-206	LSD
1999	Early county	16.1	*	18.8	1.7
2000	Early county	16.5	*	21.0	2.1
2001	Very early county	17.0	*	19.4	1.8
2001	Early county	19.5	NS	20.8	1.5
2002	Very early county	19.3	*	23.9	1.8
2002	Early county	17.5	*	19.0	1.3

^{*} Significantly different from M-104 at the 0.05 probability level.

Table 3. Days to 50 % heading for M-104, M-206 & M-202 from 1998-2002 at the Rice Experiment Station, Biggs, CA.

Year	Trial	M-104		M-206		M-202	LSD
1998	Very early Preliminary			78	1	80	2.0
1999	Early county	79	*	85	†	90	2.2
2000	Early county	71	*	75	†	78	1.5
2001	Very early county	72	*	77	†	82	2.3
2001	Early county	72	*	77	†	81	3.2
2002	Very early county	81	*	85	†	89	3.3
2002	Early county	78	*	82	Ť	88	3.1

^{*} Significantly different from M-104 at the 0.05 probability level.

18c. Exhibit C. Objective Variety Description - Rice

See attached US Plant Variety Protection form "Objective Variety Description – Rice (*Oryza sativa L*)."

[†] Significantly different from M-202 at the 0.05 probability level.

[†] Significantly different from M-202 at the 0.05 probability level.

U.S. DEPARTMENT OF AGRICULTURE PLANT VARIETY PROTECTION OFFICE, AMS, USDA NATIONAL AGRICULTURAL LIBRARY Bldg., Rm. 500 10301 BALTIMORE Blvd. BELTSVILLE, MD 20705

OBJECTIVE VARIETY DESCRIPTION RICE (Oryza sativa)

Vame	of Applic	ant(s)	Temp	orary Designation	Variety Name	
	Cali	fornia Cooperative Rice Research		98-Y-242	M-206	
		dation, Inc.				
Addres	ss (Street	and No. or R.F.D. No., City, State, and Zip Code)			FOR OFFICIAL USE	ONLY
	Rice	Experiment Station			PVPO Number	
	955	Butte City Hwy. (162)				\ .
		Box 306			2003001	80
lace t		s, CA 95917 priate number that describes the character of this variety in	tho emana-		T1	. 1 ,
orrest	onding t	o descriptors developed by IBGR-IRRI Rice Advisory Convelty more readily by describing as many characters as is pos	imittee and	i the US Rice Crop	Advisory Committee.	Breeders wil
. MA	TURITY	- Days to Heading (Seeding to 50% Heading):				
١.	South:	(Location:) at		_ kg/ha (Nitrogen rate)	
					_ kg/na (141110gen late)	
		Number of days				
÷		Days earlier than Check variety:				
		Days same as Check variety:			•	
		Days same as Check variety.				
		Days later than Check variety:				
		Maturity Class (50% heading) - South: 1 = Very earl 3 = Intermed	y (less thar iate (101 -	1 86 days) $2 = \text{Ear}$ 115) $4 = \text{Lat}$	rly (86 - 100) te (more than 115)	
3.	Califor	nia: (Location: Biggs, CA)	at	135	kg/ha (Nitrogen rate)	 r
	79	Number of days	(fall	lowed land)		
		Days earlier than Check variety: M-202				
		Days same as Check variety:				
	4	David latematican Classical M 104				
		Days later than Check variety: M-104				
	2	Maturity Class (50% heading) - California: 1 = Very earl 3 = Intermedi			ely (91 - 97) te (more than 104 days)	
CUL	M:			· · · · · · · · · · · · · · · · · · ·		-
	1	ANGLE (Degrees from Perpendicular after Flowering:				
	e e e e e e e e e e e e e e e e e e e	1 = Erect (less than 30°) 3 = Intermediate (about 45°) 7 = Spreading (more than 60° but the culms do not rest on		pen (about 60°)		

9 = Procumbent (the culm or its lower part rests on the ground surface)

LENGTH

```
9 6 6 cm (Soil level to top of extended panicle on main stem)
                       cm Shorter than Check variety: M-202
                       Length same as Check variety:
                       cm Longer than Check variety: M-104
        1
               HEIGHT CLASS:
                                       1 = Semidwarf 2 = Short
                                                                       3 = Medium
                                                                                       4 = Tail
               INTERNODE COLOR (After flowering): 1 = Green
                                                                       2 = Light Gold
                                                                                      3 = Purple lines 4 = Purple
        _3_
               STRENGTH (Lodging resistance): 1 = Strong (no lodging)
                                                                                       3 = Moderately strong (most plants leaning)
                                               5 = Intermediate (most plants lodged)
                                                                                       7 = Weak (most plants flat)
                                               9 = Very weak (all plants flat)
3. FLAG LEAF (After Heading):
       39.4
                      cm LENGTH
                                                               mm WIDTH
                                               1 2 . 2
               PUBESCENCE: 1 = Glabrous
                                               2 = Intermediate
                                                                       3 = Pubescent
        1
               LEAF ANGLE (after heading):
                                               1 = Erect
                                                               3 = Intermediate 5 = Horizontal
                                                                                               7 = Descending
        2
               BLADE COLOR:
                                       1 = Pale Green
                                                               2 = Green
                                                                               3 = Dark Green 4 = Purple tips
                                       5 = Purple margins
                                                               6 = Purple blotch 7 = Purple
               BASAL LEAF SHEATH COLOR: 1 = Green
                                                               2 = Purple lines 3 = Light purple 4 = Purple
4. LIGULE:
                      mm LENGTH (from base of collar to the tip, at late vegetative stage)
        1
              COLOR (Late vegetative state):
                                               1 = White
                                                               2 = Purple lines
                                                                                       3 = Purple
       1
              SHAPE: 1 = Acute to acuminate
                                               2 = 2-Cleft
                                                               3 = Truncate
              COLLAR COLOR (late vegetative stage): 1 = Pale green 2 = Green
                                                                                       3 = Purple
              AURICLE COLOR (late vegetative stage): 1 = Pale green
                                                                       2 = Purple
5. PANICLE:
       19.8
                      cm LENGTH
              TYPE: 1 = Compact
                                       5 = Intermediate
                                                               9 = Open
       2
              SECONDARY BRANCHING:
                                               1 = Absent
                                                               2 = Light
                                                                               3 = Heavy
                                                                                               4 = Clustering
       3
              EXSERTION (near maturity):
                                               1 = Less than 90\%
                                                                       2 = 90 - 99\%
                                                                                       3 = 100\% exserted
         2
              AXIS: 1 = Straight
                                       2 = Droopy
        1
              SHATTERING: 1 = \text{Very low (less than 1\%)}
                                                               3 = \text{Low} (1 - 5\%) 5 = \text{Moderate} (6 - 25\%)
                              7 = Moderately high (26 - 50\%)
                                                               9 = \text{High (more than 50\%)}
        3
              THRESHABILITY:
                                       1 = Difficult
                                                       2 = Intermediate
                                                                               3 = Easy
```

7. GRAIN	(Seec	i): (continue	i)											
	7_	GELATINIZATION TEMPERATURE TYPE:			1 = Hi	gh5 = I	ntermedia	te	7 = L	003	00	180		
	mylo≀ eak		e Viscosity (I ot Paste	Brabender		ed Paste		'Brea	ıkdown' 'Se	etback'				
26	69		163		26	56	*	9!	5 –	4				-
3. RESISTA	ANC	E TO LOW	TEMPERAT	URE:	······································				· · · · · · · · · · · · · · · · · · ·					
_3	3	GERMINA	ATION AND	SEEDLI	NG VIG	OR:	1 = Lc	w 2 = N	/ledium	3 = H	ligh			
	3	FLOWERI	NG (Spikele	t fertility)	:		1 = Lo	w 2 = N	⁄ledium	3 = H	ligh			
9. SEEDLI	NG V	IGOR NOT	RELATED	TO LOW	TEMPE	RATURE	:			· <u>·</u>	 -			
3	3	VIGOR: 1	= Low 2 = M	ledium	3 = H	igh					: ·			÷ "
10. BLAST 0 =	RES = Imm	ISTANCE (A	<i>Pyricularia (</i> = Resistant	<i>Oryzae</i>). 3 = M	(International oderately	ional races resistant	found u 5 = Int	nder ref ermedia	erences) ite 7 = M	oderatel	y suscepti	ble	9 = Su	sceptible
Group			ΪΒ			IC.		ID		IE	IG	IH		
Number		1 5	45	49	54	1	17	1	13	1	1	1		
Resistance			· -			_		· ·	·		9			
			ER DISEASI						· · · · · · · · · · · · · · · · · · ·				· · · · · · · · · · · · · · · · · · ·	
0=	= Imm	une I =	= Resistant	3 = M	oderately	resistant	5 = Int	ermedia	ite $7 = Mc$	oderately	y suscepti	ble	9 = Su	sceptible
	 -	Narrow Bro	own Leaf Spo	ot <i>Cercos</i> į	ora oryz	ае	7	Aggre	egate Shea	ith Spot	Rhizocton	ia oryzae	:-sativae	
+* <u></u>		Leaf Smut A	Entyloma ory	rzae				Straig	ght Head					
· · ·		Brown Leaf		nthospori olaris ory chslera o	zae)	ie	·	Kerne	el Smut		ossia horr etia barcl			
·		Leaf Scald	Gerlachia or	yzae				White	: Tip Nem	atode Ap	helencho	ides bess	eyi	
· ·	<u></u>	Hoja Blanca	a Virus		•		7	Stem	Rot <i>Sclero</i>	otium ory	yzae			
·. ———		Sheath Rot.	Sarocladium	oryzae										
 -	<u>.</u> .	Pythium Sec	edling Blight	Pythium	sp.			Bacte	rial Blight	Xantho	monas ca	mpestris]	ov. oryza	e
-	_	Sheath Spot	Rhizoctonia	oryzae			· .	Sheatl	h Blight <i>R.</i>	hizocton	ia solani			-
·		Other:			·						•			
2. INSECT	RES	STANCE:			· · · · · · · · · · · · · · · · · · ·					:	•	- 3		
	Imm		Resistant	3 = Mc	derately	resistant	5 = Inter	rmedia	te $7 = Mc$	derately	susceptil	ole	9 = Su	sceptible
	<u> </u>	Grasshopper	T	٠				Rice S	Stink Bug	Oebalus	pugnax			
. ——	_ 1	Rice Leafho	pper	٠.				Swarn	n Caterpill	lar		I.		
	_ 1	Rice Hispa					9	Rice V	Water Wee	vil <i>Lisso</i>	orhoptrus	oryzophi	lus	
· ·	_ 1	Rice Midge						Rice S	stalk Borer	Chilo p	lejadellus	7		
· · · · · · · · · · · · · · · · · · ·		Least Skippe						Sugaro	cane Borer	r Diatrae	ea saccha	ralis		

200300180

REFERENCES

- 1. C. R. Adair et al. 1972. Rice in the United States: Varieties and Production. USDA Handbook No. 289 (Rev.), 124 pp.
- 2. J. G. Atkins et al. 1967. An International Set of Rice Varieties for Differentiating Race of Pyricularia Oryzae. Phytopath. 57:297-301.
- 3. IBPGR-IRRI Rice Advisory Committee. 1980. Descriptors for Rice Oryza sativa L.). International Rice Research Institute. 21 pp.
- 4. K. C. Ling and S. H. Ou, 1969. Standardization of the International Race Numbers of Pyricularia Oryzae. Phytopath. 59:339-342.
- 5. B. D. Webb et al. 1985. Utilization Characteristics and Qualities of United States Rice. In Proceedings on Rice Grain Quality and Marketing. International Rice Research Institute (IRRI), Los Banos, Philippines. p. 25-35.

RELEASE OF 'M-206' RICE

A. Variety Name, Temporary Designation, and Genus Species

M-206

98-Y-242

Oryza sativa L.

B. Origin and Breeding History of the Variety

- 1. Genealogy: M-206 is a very early, to early, semidwarf, California glabrous, Calrose quality medium-grain (CRMG) experimental line that has been evaluated in large plot yield tests since 1998. It is a very early selection from the 1991-92 winter greenhouse cross R18087 and has the pedigree S-301/M-204. Both varieties used in the pedigree were developed by the California Cooperative Rice Research Foundation, (CCRRF) at the Rice Experiment Station (RES), Biggs, CA. S-301 is an intermediate maturity short-grain semidwarf that is no longer in production. M-204 is an early maturing Calrose medium-grain with semidwarf height. M-206 is the product of modified pedigree breeding. Early spray painted pancile selections from the 1993 F₂ nursery selected for medium grain shape were bulked 6 panicles per row in the 1993-94 Hawaiian nursery. Panicle selections grouped from the R18087 Hawaii rows were planted in 1994 individual rows. The line of descent from was 94 row 63811 via individual panicle selection to 95 row 6309 to 96 row 5136 which was bulked for 97 small plot 351 that was the seed source for entry 98-Y-242 in the RES Preliminary Yield Test. It has been tested in the University of California Cooperative Extension (UCCE) Statewide Yield Tests (99-Y-84, 00-Y-63, 01-Y-12, 01-Y-72, 02-Y-11, 02-Y-73) 1999 to 2002. Approximately 600 cwt of foundation seed is available if released.
- 2. **Selection and multiplication:** Panicle selections were first taken for headrow production in 1999 and continued through 2002. The 1999 panicles were initially purified in the 2000 headrow nursery. Year 2000 selected headrows were generation advanced in 2000-01 Hawaiian nursery. Year 2001 breeder headrows totaled 300 rows from 2000 selected headrows and their descendents from 2001-02 Hawaii nursery. The year 2001 breeder increase consisted of two components. One component was derived from bulked year 2000 headrows while the other came from the 2000-01 Hawaii nursery. These bulks were harvested separately and planted in different foundation field locations in 2002 for a total of 7.7 acres. Previous experience revealed that in certain years more outcrossing occurs in the Hawaii nursery resulting in more offtypes in either the breeder and/or foundation production fields. Two years of observations and plant counts demonstrated that 2000-01 outcrossing in Hawaii was near non-existent resulting in the two components being combined for 2002 Foundation seed. The 2001 headrow panicle selections were used to plant 400 headrows in 2002.
- 3. Variants: The breeder, foundation, and headrows were rogued multiple times and off-types, putative out-crosses, and other visual variants were removed. These included pubscent and glabrous longer medium-grains with and without red apiculus, pubscent shorter length medium-grain, slightly earlier and later maturing medium-grains, and fully awned medium grains possibly from outcrossing. In addition there were putative tetraploids (larger kernel size, delayed maturity and high floret sterility) and chlorophyll chimeras removed from the field. Total frequency of these off types was 0.0172%.

4. **Stability:** Selection, headrow, isolation, inspection, and seed production methods used by the RES for California rice varieties were followed for M-206 and no indication of problems in stability were observed in the foundation seed.

C. Description of Variety

- 1. **Botanical description:** M-206 is a medium-grain cultivated rice that is classified as Gramineae, <u>Orvza sativa L.</u>, and a temperate race japonica. It is a very early, to early, maturing semidwarf cultivar. The floret hull and leaves are glabrous except for hairs on lemma keel and leaf margins (California glabrous). Leaves are erect and green in color. The spikelet and apiculus are straw colored and sparsely awned. The appearance and amount of awns vary by years and growing environments. M-206 has a colorless, nonaromatic, and nonwaxy endosperm with a light brown pericarp.
- 2. **Objective description:** See attached USDA Plant Variety Protection Office description form Exhibit C.

D. Evidence of Variety Identity and Performance Characteristics

1. Performance characteristics: M-206 is a semidwarf, very early to early maturing, Calrose medium-grain experimental line that has been evaluated in UCCE Statewide Yield Tests since 1999 to 2002. Table 1 represents the overall evaluation from these tests plus milling, grain physical attributes, and basic starch classification characters. Comparison for these evaluation purposes were made to M-104 and M-202. Supplemental Tables 1 to 26 plus RVA graphs provide additional supporting information describing milling results, location year yield comparisions, grain physical attributes, and a more in depth quality characterization. M-103 and M-205 have been added for specific comparisions to support a more complete variety performance package. Table 1 shows that M-206 is 4 days earlier, has better lodging resistance, higher blanking resistance, improved headrice compared to M-202. M-206 has a small increase in seed weight, and size when compared to M-202 and M-104. Field observations indicate it is more stable in less than optimal field conditions, and has less nonsynchrous heading. It is not resistant to California rice blast race IG1.

Table 1. 1998 to 2002 Summary of Agronomic Characteristics for M-104, M-206, and M-202.

Character	M-104	M-206	M-202
Seedling vigor (score)	4.5	4.5	4.6
Days 50% heading	82	85*	89
Plant height (cm)	93	95	97
Lodging (%)	50	31	38
Blanking – Greenhouse (%)	13	12	25
Blanking – Davis (%)	7	12	16
Blanking – San Joaquin (%)	9	7	27
Overall blanking mean (%)	13	10	23
Stem rot (score)	5.6	5.4	5.5
Harvest moisture (%)	17.6	19.8	19.7
Yield (lb/acre @ 14%)	9087	9350*	9060
Milling (%) total	68.9	69.0	68.7
Milling (%) whole grain	63.4	66.0	64.2
Brown weight 1000 seeds (g)	24.0	24.4	22.7
Brown length (mm)	6.26	6.23	5.96
Brown width (mm)	2.72	2.75	2.80
Ratio (L/W)	2.30:1	2.27:1	2.13:1
Apparent Amylose (%)	17.2	17.4	16.2
Alkali score (1.7% KOH)	6.4	6.7	7.0

^{*}Significantly different LSD 5%.

Cold rice growing regions in Escalon and the Delta have lower temperatures along with marine air influence that generally delays maturity, reduces height and lodging, and has a potential for higher floret sterility due to lower temperatures (Table 2). M-206 behaves as a full season variety but maintains satisfactory yield performace when compared to the only commercially grown medium-grain varieties M-104 and M-103. M-206 yield performance relative to M-104 and M-103 was verified by a 2002 strip trial (Supplemental Tables 22 & 23). M-206 milling total and whole grain was improved over M-104 (Supplemental Table 3) at this location.

Table 2. 2001 to 2002 Summary of Agronomic Characteristics for M-104, M-206, M-202 and M-103 at San Joaquin.

Character	M-104	M-206	M-202	M-103
Seedling vigor (score)	4.6	4.6	4.7	4.4
Days 50% heading	93	97	98	88
Plant height (cm)	80	86	83	80
Lodging (%)	1	1	1	1
Harvest moisture (%)	21.6	23.9	24.0	19.4
Yield (lb/acre @ 14%)	8900	8590*	7880	8355

Significantly different LSD 5%.

2. Quality characteristics: Kernels of M-206 are slightly larger in size and weight compared to M-104 and M-202 (Table 1). Whole kernel grain M-206 milling %'s are improved (multi-year milling rows, plots, seed increases, and strip trials) when compared to M-104 and M-202 (Supplimental Tables 1-4). Laboratory analysis for physicochemical characteristics (apparent amylose content, protein, RVA curves, and gelatinization temperature) by the USDA Rice Quality Laboratory indicate that it fits CRMG standards. Milled rice samples of M-206, M-104, and M-202 were distributed to various California rice marketing organizations and individual rice quality evaluators in 2001-02. Responses from evaluations indicate M-206 is similar to M-104 and M-202 for various cooking and taste characters. M-206 can be commingled with other CRMG's (Supplemental Tables 25 and 26).

E. Area of Adaptation and Primary Use

M-206 is being released as a Calrose medium-grain. The data indicates it has broad adaptation in the entire rice growing region. In warm areas it will be another alternative to M-202 to allow the harvest spread for growers using M-205. Going south to the Yolo county line it can be used as an alternative or in combination with M-202. In cool production areas (Sacramento Airport, Yolo county, and areas east of Hwy 70) it can be an alternative to M-202 or be used in combination with M-202. In the coldest areas (Escalon and the Delta) it becomes a full season variety but has good blanking resistance and improved whole grain head rice over M-104.

F. Seed Maintenance

M-206 was developed and is owned by CCRRF. It is being proposed for joint release by CCRRF, the California Agriculture Experiment Station (CAES), and the United States Department of Agriculture—Agricultural Service (USDA-ARS). Breeder and foundation seed of M-206 will be maintained by CCRRF at the Rice Experiment Station, P. O. Box 306, 955 Butte City Highway, Biggs, CA 95917-0306. Foundation seed will be planted to produce registered seed and registered seed will be planted to produce certified seed. Headrow seed will be produced as needed in the foundation seed field to minimize outcrossing and insure parity. Foundation seed can be used to produce foundation seed when necessary. The various classes of seed will be produced in California. Application will be made for M-206 under the U. S. Plant Variety Protection, Title V option, and a utility patent also will be applied for by CCRRF.

REPRODUCE LOCALLY. Include form number and edition date on a	I reproducti	
AIGH DENAKT MENT OF AGRICUIT LIBE	in reproductions.	FORM APPROVED - OMB No. 0581-00
AGRICULTURAL MARKETING SERVICE	Application is required in order to de	termine if a plant various protoction
EXHIBIT E		
STATEMENT OF THE BASIS OF OWNERSHIP	confidential until the certificate is issu	ued (7 U.S.C. 2426).
1. NAME OF APPLICANT(S) California Cooperative Rice Research	2. TEMPORARY DESIGNATION	3. VARIETY NAME
	OR EXPERIMENTAL NUMBER	J. VARIE! Y NAIVIE
Foundation, Inc.	98-Y-242	M-206
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP, and Country)	5. TELEPHONE (Include area code)	I G FAY
Rice Experiment Station	, and the second	6. FAX (Include area code)
P.O. Box 306	(530)868-5481	(530)868–1730
Biggs, CA 95917-0306	7. PVPO NUMBER	
		0180
8. Does the applicant own all rights to the variety? Mark an "X" in the	appropriate block. If no plant	
	Thu, please expl	X YES NO
•		
9. is the applicant (individual or company) a LLC Avi		
9. is the applicant (individual or company) a U.S. National or a U.S. ba	ased company? If no, give name of co	ountry YES NO
		X 123 NO
10. Is the applicant the original owner?	· ·	•
NO Is the applicant the original owner?	If no, please answer one of the foll	owing:
a If the original value of		
a. If the original rights to variety were owned by individual(s), is (a	re) the original owner(s) a U.S. Nationa	l(s)?
T YES T NO	If no, give name of country	
	•	
b. If the original rights to variety were owned by a company(les), i	S (are) the original gumos(s) = 11.0	
		ed company?
YES NO	If no, give name of country	
· · · · · · · · · · · · · · · · · · ·		
11. Additional explanation on ownership (If needed, use the reverse for	Overs angels	
M-206 was developed by Dr. Carl W. Johnson		r employee by the
	_	r emproyee by the
California Cooperative Rice Research Four	ndation, inc. (CCRRF)	
PLEASE NOTE:		
Plant variety protection can only be afforded to the owners (not licensee		
1 If the date to the	s) who meet the following criteria:	
 If the rights to the variety are owned by the original breeder, that pers national of a country which affords similar protection to nationals of th 	on must be a U.S. national, national of	a UPOV member country, on
If the rights to the variety are owned by the company which employed nationals of a UPOV member country, or owned by nationals of a country.	the original breeder(s), the company m	nist holl S. based and t
nationals of a UPOV member country, or owned by nationals of a cou- genus and species.	ntry which affords similar protection to	nationals of the U.S. for the same
•	the state of the s	
3. If the applicant is an owner who is not the original owner, both the original owner, both the original owner.	ginal owner and the applicant must mee	t one of the above criteria
The original breeder/owner may be the individual or company who direct Act for definitions.	ed the final breeding. See Seeting 44	
ACLIOI GENNIONS.		a)(2) of the Plant Variety Protection
According to the Paragraph Code and		
According to the Paperwork Reduction Act of 1995, an agency may not conduct or sponsor, and control number. The valid OMB control number for this information collection is 0581-0055. The response, including the time for reviewing the instructions, searching existing data sources, gath	a person is not required to respond to a collection of	information unless it displays a valid OMB
to remain the instructions, searching existing data sources, gathe	Sting and maintaining the state of the Conscious	is esumated to average 6 minutes per
The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs on the ba- ind marital or familial status. (Not all prohibited bases apply to all programs). Persons with disab- judiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).	sis of race, color, national origin, gender, religion, ad	e, disability, political beliefs, sexual orientation
reducible, erc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD).	wild require alternative means for communical	ion of program information (braille, large print,